

CERTHIA CYANEA.



CHARACTER GENERICUS.

Rostrum arcuatum, tenue, subtrigonum, acutum.

Lingua acuta.

Pedes ambulatorii.

Lin. Syst. Nat. p. 184.

CHARACTER SPECIFICUS, &c.

CERTHIA CYANEA, vertice beryllino, alis caudaque nigris, tectricibus inferioribus flavis.

CERTHIA cærulea, fascia oculari humeris alis caudaque nigris, pedibus rubris.

Lin. Syst. Nat. p. 188.

CERTHIA cærulea, fascia oculari, humeris, alis caudaque nigris, pedibus fuscis.

Lin. Syst. Nat. Gmel. p. 483.

Certhiæ familiari æqualis prope est magnitudine avis quam repræsentat tabula. Generatur in America Australi, in Cayana autem præcipue.



THE
CYANEAN CREEPER.



GENERIC CHARACTER.

Bill slender, incurvated, sharp-pointed.
Tongue differing in shape in the different species.
Legs moderately stout.
Toes three before; one behind: back-toe large;
claws hooked and long.

SPECIFIC CHARACTER, &c.

Blue CREEPER, with the wings and tail black;
the crown beryl-coloured; the under wing-
coverts yellow.

GUIT-GUIT noir et bleu.

Buff. ois. 5. p. 529.

GRIMPEREAU du Bresil.

Pl. Enl. 83. f. 2.

Black-and-blue CREEPER.

Edw. pl. 264. f. 1.

The bird represented on the plate is an inhabitant of South America, and is principally found in the province of Cayenne. Its size is nearly that of the common English Creeper.